arXiv:1209.2285v1 [quant-ph] 11 Sep 2012

Optimal transfer of an unknown state via a bipartite operation

Yang Liu,[1] Yu Guo,[1, 2] and D. L. Zhou[1]

[1] *Beijing National Laboratory for Condensed Matter Physics,*
and Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China
[2] *School of Physics and Electronic Science, Changsha University of Science and Technology, Changsha 410114, China*

A fundamental task in quantum information science is to transfer an unknown state from particle A to particle B (often in remote space locations) by using a bipartite quantum operation $\mathcal{E}^{AB}$. We suggest the power of $\mathcal{E}^{AB}$ for quantum state transfer (QST) to be the maximal average probability of QST over the initial states of particle B and the identifications of the state vectors between A and B. We find the QST power of a bipartite quantum operations satisfies four desired properties between two d-dimensional Hilbert spaces. When A and B are qubits, the analytical expressions of the QST power is given. In particular, we obtain the exact results of the QST power for a general two-qubit unitary transformation.

PACS numbers: 03.67.-a, 03.65.-w

I. INTRODUCTION

A fundamental task in quantum information science is to transfer an unknown internal quantum state of a particle from one location $\mathcal{A}$ to another location $\mathcal{B}$. A direct method is to mechanically move the particle from $\mathcal{A}$ to $\mathcal{B}$ while keeping the internal state invariant. A more sophisticated way is quantum state teleportation [1], where the unknown state is teleported with the aid of a pair of particles in a Bell state and 2 bits of classical communications. The third way is to transfer the state via a two-particle quantum operation $\mathcal{E}^{AB}$, which can be realized by linking two nodes A and B to a quantum network, e.g., a quantum wire (a one-dimensional chain of particles with interactions) [2–4]. Here the node A, located in $\mathcal{A}$, is the particle with the unknown state to be transferred, and the node B, located in $\mathcal{B}$, is the particle as the state receiver.

The aim of Refs. [3–9] is to achieve perfect quantum state transfer by optimizing the quantum network. In addition, the capacity of quantum state transfer to characterize the non-locality of a bipartite unitary transformation is studied in Refs. [10, 11]. Here we will solve another related question: For a given two-particle quantum operation $\mathcal{E}^{AB}$, what is the maximal average probability for quantum state transfer? This maximal probability reflects the power of quantum state transfer of the bipartite operation $\mathcal{E}^{AB}$.

The approach we will adopt is similar to that in the power of entanglement generation for a local unitary gate [12–14]. Here we want to emphasize that the bipartite quantum operations, including non-unitary gates, are necessary to be considered for quantum state transfer.

The article is organized as follows. In Sec. II, we introduce the basic formula of the power of quantum state transfer, and four basic properties for the QST power are proved. In Sec. III, we give the analytical results of two-qubit operations. In particular, an exact result of the QST power for any two-qubit unitary transformation is given. Finally we present some discussions and a brief summary.

II. GENERAL RESULTS

A. The power of QST

In this section, we will give a proper quantity to measure the power of QST for a bipartite quantum operation.

We consider two particles A and B, whose Hilbert space is $\mathcal{H}^{AB} = \mathcal{H}^A \otimes \mathcal{H}^B$ with $\dim \mathcal{H}^A = \dim \mathcal{H}^B = d$. In other words, particles A and B are two qudits. Initially particle A is prepared in an unknown state $|\psi^A\rangle$, and particle B is in some given state $|\xi^B\rangle$. For convenience, we take $|\psi^A\rangle = R|0^A\rangle$ with $|0^A\rangle$ being any given state and $R \in SU(d)$. In general, there are many choices of R for given $|\psi^A\rangle$ and $|0^A\rangle$, which does not affect the following formulations. After performing a bipartite quantum operation $\mathcal{E}^{AB}$, we need to estimate to which degree the unknown state $|\psi^A\rangle$ being transferred to partilce B. This process of QST is depicted in Fig. 1.



FIG. 1. The process of quantum state transfer using a bipartite quantum operation $\mathcal{E}^{AB}$. We maximize the probability of QST over particle B's initial states $|\xi^B\rangle$ and the unitary transformations S^B.

For a given bipartite quantum operations, we may improve the QST by controlling two elements. On one hand, the QST power of $\mathcal{E}^{AB}$ depends on the input state of particle B $|\xi^B\rangle$. We can improve the QST by preparing particle B in a suitable initial state. On the other hand, notice that if the final state is $S^B\left|\psi^B\right\rangle$, where S^B is a unitary transformation on particle B independent of the unknown state $|\psi^A\rangle$, the unkown state will be regarded as being perfectly transferred. Different choices of S^B implies different identifications of the bases between $\mathcal{H}^A$ and $\mathcal{H}^B$. Hence we can improve the QST of $\mathcal{E}^{AB}$ by adjusting S^B.

To give a measure to characterize the power of QST for $\mathcal{E}^{AB}$, we need to optimize over $|\xi^B\rangle$ and S^B. Therefore the QST power for a quantum operation $\mathcal{E}^{AB}$ is defined as

$$\mathcal{P}_{QST}\left(\mathcal{E}^{AB}\right)=\max_{S^B,|\xi^B\rangle}\int d\mu\left(R\right)P(R;\mathcal{E}^{AB};S^B,|\xi^B\rangle) \tag{1}$$

where

$$P(R;\mathcal{E}^{AB};S^B,|\xi^B\rangle)=\mathrm{Tr}\left(\mathcal{E}^{AB}\mathcal{R}^A\left(P_0^AP_\xi^B\right)\mathcal{S}^B\mathcal{R}^B(P_0^B)\right) \tag{2}$$

with $P_0^A=|0^A\rangle\langle0^A|$, $P_\xi^B=|\xi^B\rangle\langle\xi^B|$, $P_0^B=|0^B\rangle\langle0^B|$, $\mathcal{R}$ and $\mathcal{S}$ are the local quantum operations corresponding to the unitary transformations R and S, and $d\mu\left(R\right)$ being the Haar measure on $SU\left(d\right)$. Because the Harr measure is right-invariant, the power of QST is independent of the choice of $|0\rangle$. Since we have no reasons to give different probabilities to two sets of states connecting by a unitary transformation in the average over the unknown states, the Harr measure is a natural choice for the average.

For a given initial state $R^A|0\rangle\otimes|\xi^B\rangle$ and a given identification between $\mathcal{H}^A$ and $\mathcal{H}^B$ specified by S^B, $P(R;\mathcal{E}^{AB};S^B,|\xi^B\rangle$ is the probability of QST for the state $R|0\rangle$ by $\mathcal{E}^{AB}$. Further more, the power of QST for $\mathcal{E}^{AB}$ is the maximal average probability for particle B in the unkown state after the action of $\mathcal{E}^{AB}$.

B. Properties of the QST power

In the above subsection, we give a definition of the QST power for a bipartite quantum operation, and give its physical interpretation. Here we will prove that the QST power $\mathcal{P}_{QST}\left(\mathcal{E}^{AB}\right)$ has the following desired properties.

Property (i). The QST power is invariant under local unitary transformations.

Let X^A, Y^B, U^A, V^B be local unitary transformations, and let $\mathcal{X}^A$, $\mathcal{Y}^B$, $\mathcal{U}^A$, $\mathcal{V}^B$ be the corresponding local quantum operations. Then

$$\begin{aligned}
&\mathcal{P}_{QST}\left(\mathcal{X}^A\mathcal{Y}^B\mathcal{E}^{AB}\mathcal{U}^A\mathcal{V}^B\right)\\
&=\max_{S^B,|\xi^B\rangle}\int d\mu(R)P(R;\mathcal{X}^A\mathcal{Y}^B\mathcal{E}^{AB}\mathcal{U}^A\mathcal{V}^B;S^B,|\xi^B\rangle)\\
&=\max_{S^B,|\xi^B\rangle}\int d\mu(R)P(UR;\mathcal{E}^{AB};Y^{B\dagger}S^BV^{B\dagger},V^B|\xi^B\rangle)\\
&=\max_{S^B,|\xi^B\rangle}\int d\mu(R)P(R;\mathcal{E}^{AB};S^B,|\xi^B\rangle)\\
&=\mathcal{P}_{QST}\left(\mathcal{E}^{AB}\right),
\end{aligned} \tag{3}$$

where we have used Eq. (A2) in the third line of the above equation.

Property (i) shows the power of a bipartite quantum operation characterizes its nonlocal property [14].

Property (ii). The range of the QST power is in the period $\left[\frac{1}{d},1\right]$, i.e.,

$$\frac{1}{d}\le\mathcal{P}_{QST}\left(\mathcal{E}^{AB}\right)\le1. \tag{4}$$

This can be proved as follows. Because

$$\begin{aligned}
&\mathcal{P}_{QST}\left(\mathcal{E}^{AB}\right)\\
&=\max_{S^B,|\xi^B\rangle}\int d\mu\left(R\right)P(R;\mathcal{E}^{AB};S^B,|\xi^B\rangle)\\
&\le\max_{S^B,|\xi^B\rangle}\int d\mu\left(R\right)1=1.
\end{aligned}$$

In addition, if $\mathcal{P}_{QST}\left(\mathcal{E}^{AB}\right)<\frac{1}{d}$, then

$$\begin{aligned}
&\int d\mu\left(S\right)\int d\mu\left(R\right)P(R;\mathcal{E}^{AB};S^B,|\xi^B\rangle)\\
&\le\int d\mu\left(S\right)\mathcal{P}_{QST}\left(\mathcal{E}^{AB}\right)<\frac{1}{d}.
\end{aligned}$$

However,

$$\begin{aligned}
&\int d\mu\left(S\right)\int d\mu\left(R\right)P(R;\mathcal{E}^{AB};S^B,|\xi^B\rangle)\\
&=\int d\mu\left(R\right)\mathrm{Tr}\left(\mathcal{E}^{AB}\left(R^AP_0^AR^{A\dagger}P_\xi^B\right)\frac{I^B}{d}\right)\\
&=\int d\mu\left(R\right)\frac{1}{d}=\frac{1}{d}.
\end{aligned}$$

In the second line, we used the lemma (A3). This result contradicts with the above inequality. Therefore $\frac{1}{d}\le\mathcal{P}_{QST}\left(\mathcal{E}^{AB}\right)$.

Notice that the lower bound $1/d$ is the same as in the case of transferring a classical discrete variable with d different states.

Property (iii). The QST power of a local operation $\mathcal{E}^A\otimes\mathcal{E}^B$ is $\frac{1}{d}$.

$$\begin{aligned}
&\mathcal{P}_{QST}\left(\mathcal{E}^A\otimes\mathcal{E}^B\right)\\
&=\max_{S^B,|\xi^B\rangle}\int d\mu\left(R\right)P(R;\mathcal{E}^A\mathcal{E}^B;S^B,|\xi^B\rangle)\\
&=\max_{S^B,|\xi^B\rangle}\int d\mu\left(R\right)\mathrm{Tr}\left(\mathcal{E}^B\left(P_\xi^B\right)S^BR^BP_0^BR^{B\dagger}S^{B\dagger}\right)\\
&=\max_{|\xi^B\rangle}\int d\mu\left(R\right)\mathrm{Tr}\left(\mathcal{E}^B\left(P_\xi^B\right)R^BP_0^BR^{B\dagger}\right)\\
&=\max_{|\xi^B\rangle}\mathrm{Tr}\left(\mathcal{E}^B\left(P_\xi^B\right)\frac{I^B}{d}\right)\\
&=\frac{1}{d}.
\end{aligned}$$

In the fifth line of the above equation, we use the lemma (A3).

As expected, a local operation has the lowest power in transferring an unknown state.

Property (iv). The QST power of the swapping gate $SWAP^{AB}$ is 1.

$$\begin{aligned}
&\mathcal{P}_{QST}\left(SWAP^{AB}\right)\\
&= \max_{S^B,|\xi^B\rangle}\int d\mu\left(R\right)P(R;SWAP^{AB};S^B,|\xi^B\rangle)\\
&= \max_{S^B,|\xi^B\rangle}\int d\mu\left(R\right)\operatorname{Tr}\left(R^B P_0^B R^{B\dagger}P_\xi^A S^B R^B P_0^B R^{B\dagger}S^{B\dagger}\right)\\
&= \max_{S^B}\int d\mu\left(R\right)\operatorname{Tr}\left(R^B P_0^B R^{B\dagger}S^B R^B P_0^B R^{B\dagger}S^{B\dagger}\right)\\
&= \int d\mu\left(R\right)\operatorname{Tr}\left(R^B P_0^B R^{B\dagger}R^B P_0^B R^{B\dagger}\right)\\
&= 1.
\end{aligned}$$

This result is reasonable because the unknown state is swapped, i.e., perfectly transferred.

III. ANALYTICAL RESULTS OF THE QUBIT CASE

Since the QST power is defined as an optimization problem over a state and a unitary transformation in a d-dimensional Hilbert space, the explicit calculations of the QST power of $\mathcal{E}^{AB}$ when A and B are two qudits, in general, are complex. In this section, we will give an analytical result on the QST power for an arbitrary two-qubit quantum operation $\mathcal{E}^{AB}$, which makes the numerical calculation of the QST power becomes accessible. In particular, we further obtain the exact result of the QST power for any two-qubit unitary transformation.

For the qubit case, let $P_0 = \frac{I+\sigma_z}{2}$. $\forall R \in SU(2)$, we can find a coordinate frame $\{\vec{R}_x, \vec{R}_y, \vec{R}_z\}$ to characterize it. The base vector of the coordinate frame is defined by $R\sigma_n R^\dagger = \vec{R}_n\cdot\vec{\sigma}$. Then the component of the bases vector $R_n^m = \frac{\operatorname{Tr}\left(R\sigma_n R^\dagger\sigma_m\right)}{2}$. The initial state of particle B $P_\xi^B = \frac{I^B+\vec{T}\cdot\vec{\sigma}^B}{2}$, where $\vec{T}$ is the Bloch vector for the state $|\xi^B\rangle$.

The QST power of $\mathcal{E}^{AB}$ is

$$\mathcal{P}_{QST}\left(\mathcal{E}^{AB}\right) = \frac{1}{2} + \frac{\max\limits_{S,T}\left(\sum\limits_{l,n}\mathcal{E}_{0n}^l S_l^n + \sum\limits_{l,m,n}\mathcal{E}_{mn}^l T^m S_l^n\right)}{24}, \tag{5}$$

where

$$\begin{aligned}
\mathcal{E}_{0n}^l &= \operatorname{Tr}\left(\mathcal{E}^{AB}\left(\sigma_l^A I^B\right)\sigma_n^B\right),\\
\mathcal{E}_{mn}^l &= \operatorname{Tr}\left(\mathcal{E}^{AB}\left(\sigma_l^A\sigma_m^B\right)\sigma_n^B\right),
\end{aligned}$$

and $\{S_n\}$ is the basis vectors of the coordinate frame defined by the unitary transformation S.

To derive Eq. (5), we used the following Harr average values on $SU(2)$:

$$\begin{aligned}
\langle R_z^m\rangle &= 0,\\
\langle R_z^m R_z^n\rangle &= \frac{1}{3}\delta_{mn},
\end{aligned}$$

whose detailed proofs are given in the appendix B.

Let us demonstrate the power of Eq. (5) with calculating the QST power of the CNOT gate. A direct calculation gives

$$\mathcal{P}_{QST}\left(CNOT^{AB}\right) = \frac{1}{2} + \frac{\max_{S,T}\left(T^x S_z^x + T^y S_z^y\right)}{6} = \frac{2}{3}.$$

It is worthy to point out that Eq. (5) can be used as the foundation for numerical calculations of the QST power for arbitrary two-qubit quantum operation. For example, it may find applications in the process of QST along a quantum wire [3–5].

A. Exact result on QST power for arbitrary two-qubit unitary transformations

In this subsection, we will apply Eq. (5) to the case when $\mathcal{E}^{AB}$ is a two-qubit unitary transformation. In this case, the exact result of the QST power will be obtained.

First notice that a general unitary transformation for two qubits can be written as

$$U^{AB} = U^A U^B U_d^{AB} V^B V^A,$$

where

$$U_d^{AB} = e^{-\frac{i}{2}\left(\sum_m d_m\sigma_m^A\sigma_m^B\right)}$$

with $|d_z| \le d_y \le d_x \le \frac{\pi}{2}$ [13, 15]. Because the QST power is invariant under local unitary transformations, it is sufficient to study the unitary transformation U_d^{AB}.

Through a complex but direct calculation, we arrives at

$$\mathcal{P}_{QST}\left(U^{AB}\right) = \frac{1}{2} + \frac{\max_{S,T} f}{6},$$

where

$$\begin{aligned}
f =&\sin d_y\sin d_z S_x^x + \sin d_z\sin d_x S_y^y + \sin d_x\sin d_y S_z^z\\
&+ \cos d_y T^y\sin d_x S_x^z + \cos d_z T^z\sin d_y S_y^x + \cos d_x T^x\sin d_z S_z^y\\
&- \cos d_z T^z\sin d_x S_x^y - \cos d_x T^x\sin d_y S_y^z - \cos d_y T^y\sin d_z S_z^x.
\end{aligned}$$

Notice that

$$f\left(-\vec{S}_x, -\vec{S}_y, \vec{S}_z, -\vec{T}|d_z \le 0\right) = f\left(\vec{S}_x, \vec{S}_y, \vec{S}_z, \vec{T}|d_z \ge 0\right).$$

Hence $\max f\left(d_z\right) = \max f\left(-d_z\right)$. Therefore we only need to study the case when $d_z \ge 0$.

We obtain the exact result on the maximization of f:

$$\max_{S,T} f = \sin d_x + \sin d_y + \sin d_x\sin d_y. \tag{6}$$

The proof of Eq. (6) can be found in the appendix C. Therefore the power of a two-qubit unitary transformation is

$$\mathcal{P}_{QST}\left(U^{AB}\right) = \frac{1}{2} + \frac{\sin d_x + \sin d_y + \sin d_x\sin d_y}{6}. \tag{7}$$

A remarkable feature in the QST power of U^{AB} is that it is independent of the parameter d_z. For the CNOT gate, $d_x = \pi/2$ and $d_y = d_z = 0$, so its QST power is $2/3$, which is the same as calculated above. To make a perfect QST, we require that $d_x = d_y = \pi/2$. To make the QST power lowest, we get $d_x = d_y = 0$, where U^{AB} becomes a local unitary transformation.

IV. DISCUSSIONS AND SUMMARY

In Refs. [10, 11], the nonlocal properties of a bipartite gate are classified according their capacities in transmitting classical or quantum bits of information. Here we suggest the QST power to characterize the capacity to transmit quantum state. In parallel, our method can be generalized to classical state transfer or quantum state swapping.

Technically, we prove that the QST power of a local bipartite quantum operation is $1/d$. However we don't know whether the QST power of a bipartite quantum operation is $1/d$ implies that the bipartite quantum operation is local. In addition, we give a lengthy proof of the maximization, Eq. (6), in Appendix C. Does there exist some simpler proof of Eq. (6)?

In summary, we suggest the QST power of a bipartite quantum operation as the maximal average probability of QST using the bipartite quantum operation. Four basic properties of the QST power of a bipartite quantum operation are proved. Then we obtain the analytical result of the QST power for any two-qubit quantum operation, which may be used as the foundation for numerical calculations of the QST power. The exact result of the QST power for arbitrary two-qubit unitary transformation is obtained. We hope that our work present an alternative method to characterize the non-locality of a bipartite quantum operation.

ACKNOWLEDGMENTS

We thank S.L. Luo, Z.W. Zhou, and C.P. Sun for helpful discussions. This work is supported by NSF of China (Grant Nos. 10975181 and 11175247) and NKBRSF of China (Grant No. 2012CB922104).

Appendix A: Harr measure

Notice that the Harr measure satisfies two useful properties [16]:

i) It is normalized.

$$\int d\mu(R) 1 = 1. \tag{A1}$$

ii) It is left-invariant and right-invariant. $\forall S \in SU(d)$,

$$\int d\mu(R) f(R) = \int d\mu(R) f(SR) = \int d\mu(R) f(RS). \tag{A2}$$

Lemma 1. *$\forall |0\rangle \in \mathcal{H}$ and $R \in SU(d)$, we have*

$$\int d\mu(R) R P_0 R^\dagger = \frac{I}{d}. \tag{A3}$$

This can be proved as follows. Firstly we take a complete normal orthogonal bases of $\mathcal{H}$, denoted as $\{|n\rangle,\ n \in \{0, 1, \cdots, d-1\}\}$. Because $\forall n$ there exists a unitary transformation S_n such that $|n\rangle = S_n|0\rangle$, Eq. (A2) gives $\int d\mu(R) R P_n R^\dagger = \int d\mu(R) R P_0 R^\dagger$. Therefore $\int d\mu(R) R P_0 R^\dagger = \int d\mu(R) R \frac{\sum_n P_n}{d} R^\dagger = \frac{I}{d}$.

Appendix B: Harr average on $SU(2)$

Proposition 1. *$\forall m, n \in \{x, y, z\}$, we have*

$$\langle R_z^m \rangle = 0,$$

$$\langle R_z^m R_z^n \rangle = \frac{1}{3}\delta_{mn}.$$

We can prove the above result by a direct calculation. Here we present an alternative approach as follows.

$\forall m \in \{x, y, z\}$, $\exists n \neq m$,

$$\begin{aligned}
\langle R_z^m \rangle &= \int d\mu\left(R\right) \frac{\mathrm{Tr}\left(R\sigma_z R^\dagger \sigma_m\right)}{2} \\
&= \int d\mu\left(R\right) \frac{\mathrm{Tr}\left(\sigma^n R\sigma_z R^\dagger \sigma_n^\dagger \sigma_m\right)}{2} \\
&= -\int d\mu\left(R\right) \frac{\mathrm{Tr}\left(R\sigma_z R^\dagger \sigma_m\right)}{2} \\
&= -\left\langle R_z^m \right\rangle.
\end{aligned}$$

Therefore $\langle R_z^m \rangle = 0$.

When $m \neq n$,

$$\begin{aligned}
&\langle R_z^m R_z^n \rangle = \int d\mu\left(R\right) \frac{\mathrm{Tr}\left(R^A \sigma_z^A R^{A\dagger} \sigma_m^A R^B \sigma_z^B R^{B\dagger} \sigma_n^B\right)}{4} \\
&= \int d\mu\left(R\right) \frac{\mathrm{Tr}\left(\sigma_m^A R^A \sigma_z^A R^{A\dagger} \sigma_m^{A\dagger} \sigma_m^A \sigma_m^B R^B \sigma_z^B R^{B\dagger} \sigma_m^{B\dagger} \sigma_n^{B\dagger}\right)}{4} \\
&= -\int d\mu\left(R\right) \frac{\mathrm{Tr}\left(R^A \sigma_z^A R^{A\dagger} \sigma_m^A R^B \sigma_z^B R^{B\dagger} \sigma_n^B\right)}{4} \\
&= -\left\langle R_z^m R_z^n \right\rangle.
\end{aligned}$$

Hence, $\langle R_z^m R_z^n \rangle = 0$ if $m \neq n$.

When $m \neq n$, $\exists$ a unitary transformation H, such that $H^\dagger \sigma_m H = \sigma_n$. Hence

$$\begin{aligned}
&\langle R_z^m R_z^m \rangle = \int d\mu\left(R\right) \frac{\mathrm{Tr}\left(R^A \sigma_z^A R^{A\dagger} \sigma_m^A R^B \sigma_z^B R^{B\dagger} \sigma_m^B\right)}{4} \\
&= \int d\mu\left(R\right) \frac{\mathrm{Tr}\left(H^A R^A \sigma_z^A R^{A\dagger} H^{A\dagger} \sigma_m^A H^B R^B \sigma_z^B R^{B\dagger} H^{B\dagger} \sigma_m^{B\dagger}\right)}{4} \\
&= \int d\mu\left(R\right) \frac{\mathrm{Tr}\left(R^A \sigma_z^A R^{A\dagger} \sigma_n^A R^B \sigma_z^B R^{B\dagger} \sigma_n^B\right)}{4} \\
&= \left\langle R_z^n R_z^n \right\rangle.
\end{aligned}$$

Because $\sum_m \left(R_z^m\right)^2 = 1$, we have $\langle R_z^m R_z^m \rangle = \frac{1}{3}$.

Appendix C: Maximization of the function f

In this appendix, we will prove Eq. (6). First, we divide f into two parts:

$$f = f_1 + f_2,$$

where

$$\begin{aligned} f_1 &= \sin d_y \sin d_z S_x^x + \sin d_z \sin d_x S_y^y + \sin d_x \sin d_y S_z^z, \\ f_2 &= \lambda a + \mu b + \gamma c + \eta g + \nu p + \xi q \end{aligned}$$

with $\lambda = T^z S_x^y$, $\mu = -T^z S_y^x$, $\gamma = -T^y S_x^z$, $\eta = T^y S_z^x$, $\nu = T^x S_y^z$, $\xi = -T^x S_z^y$, and $a = \cos d_z \sin d_x$, $b = \cos d_z \sin d_y$, $c = \cos d_y \sin d_x$, $g = \cos d_y \sin d_z$, $p = \cos d_x \sin d_y$, $q = \cos d_x \sin d_z$.

Because $\vec{T}$ is a unit vector, and $\{\vec{S}_x, \vec{S}_y, \vec{S}_z\}$ is a right-handed coordinate frame, they are parameterized as follows.

$$T^x = \sin\alpha \cos\beta, \tag{C1}$$

$$T^y = \sin\alpha \sin\beta, \tag{C2}$$

$$T^z = \cos\alpha, \tag{C3}$$

$$S_x^x = (\sin^2\phi + \cos^2\theta \cos^2\phi)\cos\omega + \cos^2\phi \sin^2\theta, \tag{C4}$$

$$\begin{aligned} S_x^y &= \cos\phi \sin\phi + \cos\theta \sin\omega - \cos\phi \cos^2\theta \sin\phi \\ &\quad - \cos\omega \cos\phi \sin\phi + \cos\omega \cos\phi \cos^2\theta \sin\phi, \end{aligned} \tag{C5}$$

$$S_x^z = \sin\theta(\cos\phi \cos\theta(1-\cos\omega) - \sin\omega \sin\phi), \tag{C6}$$

$$\begin{aligned} S_y^x &= \cos\phi \sin\phi - \cos\theta \sin\omega - \cos\phi \cos^2\theta \sin\phi \\ &\quad - \cos\omega \cos\phi \sin\phi + \cos\omega \cos\phi \cos^2\theta \sin\phi, \end{aligned} \tag{C7}$$

$$S_y^y = (\cos^2\phi + \cos^2\theta \sin^2\phi)\cos\omega + \sin^2\theta \sin^2\phi, \tag{C8}$$

$$S_y^z = \sin\theta(\cos\phi \sin\omega + (1-\cos\omega)\cos\theta \sin\phi), \tag{C9}$$

$$S_z^x = \sin\theta(\sin\omega \sin\phi + (1-\cos\omega)\cos\phi \cos\theta), \tag{C10}$$

$$S_z^y = \sin\theta((1-\cos\omega)\cos\theta \sin\phi - \cos\phi \sin\omega), \tag{C11}$$

$$S_z^z = \cos^2\theta + \cos\omega \sin^2\theta, \tag{C12}$$

where $0 \leq \alpha, \theta, \omega \leq \pi$, $0 \leq \beta, \phi \leq 2\pi$.

1. Analysis of f_1

Using Eq.(C4), Eq.(C8), Eq.(C12), we can rewrite f_1 as

$$\begin{aligned} f_1 &= \sin d_y \sin d_z S_x^x + \sin d_z \sin d_x S_y^y + \sin d_x \sin d_y S_z^z \\ &= (1-\cos\omega) M \cos^2\theta + \sin d_x \sin d_y \cos\omega \\ &\quad + \sin d_y \sin d_z \left(\cos^2\phi + \cos\omega \sin^2\phi\right) \\ &\quad + \sin d_x \sin d_z \left(\sin^2\phi + \cos\omega \cos^2\phi\right). \end{aligned}$$

Because

$$\begin{aligned} M &= \sin d_x \sin d_y - \sin d_y \sin d_z \cos^2\phi - \sin d_x \sin d_z \sin^2\phi \\ &\geq \sin d_x \sin d_y - \sin d_y \sin d_z \cos^2\phi - \sin d_x \sin d_z \sin^2\phi \\ &\geq \sin d_x \sin d_y \left(1 - \cos^2\phi - \sin^2\phi\right) \\ &= 0 \end{aligned}$$

and $1 - \cos\omega \geq 0$,

$$\begin{aligned} f_1 &\leq f_1(\theta = 0,\, \phi,\, \omega) = f_1(\theta = \pi,\, \phi,\, \omega) \\ &= \sin d_x \sin d_y + \cos\omega \left(\sin d_x + \sin d_y\right) \sin d_z \\ &\equiv f_{1max}. \end{aligned}$$

2. Analysis of f_2

From $0 \leq d_z \leq d_y \leq d_x \leq \frac{\pi}{2}$, we can get the inequality:

$$a \geq \left\{ \begin{array}{c} b \\ c \end{array} \right\} \geq \left\{ \begin{array}{c} g \\ p \end{array} \right\} \geq q.$$

We also have the relations

$$\begin{aligned} &\lambda + \mu + \gamma + \eta + \nu + \xi \\ &= -2\left(\cos\alpha \cos\theta + \cos[\beta - \phi] \sin\alpha \sin\theta\right) \sin\omega \\ &= -2\sqrt{\cos^2\theta + \cos^2[\beta - \phi] \sin^2\theta}\, \sin(\alpha + \varphi) \sin\omega \\ &\leq 2\left(\sqrt{1 + (\cos^2[\beta - \phi] - 1)\sin^2[\theta]}\right) \sin\omega \\ &\leq 2\sin\omega \\ &\leq 2, \end{aligned} \tag{C13}$$

where $\tan\varphi = \frac{1}{\cos(\beta - \phi)\tan(\theta)}$. The equality holds when $\beta = \phi$, $\alpha - \theta = \pm\pi$, $\omega = \frac{\pi}{2}$ or $\beta - \phi = \pm\pi$, $\alpha + \theta = \pi$, $\omega = \frac{\pi}{2}$.

$$\begin{aligned} &\lambda + \mu + \gamma + \eta \\ &= 2(\cos\alpha \cos\theta + \sin\alpha \sin\beta \sin\theta \sin\phi) \sin\omega \\ &\leq 2\sin\omega \\ &\leq 2, \end{aligned} \tag{C14}$$

$$\lambda + \gamma = T^z S_x^y - T^y S_x^z = \left(\vec{S_x} \times \vec{T}\right)_x \leq 1, \tag{C15}$$

$$\nu + \mu = T^x S_y^z - T^z S_y^x = \left(\vec{S_y} \times \vec{T}\right)_y \leq 1, \tag{C16}$$

Proposition 2. *$f_2 \leq a + b$.*

To see this, we consider

$$a + b - f_2 = a + b - (\lambda a + \mu b + \gamma c + \eta g + \nu p + \xi q)$$

1) If $-\eta g - \xi q \geq 0$,

$$\begin{aligned} &a + b - f_2 \\ &= (1-\lambda)\, a + (1-\mu)\, b - \gamma c - \nu p - \eta g - \xi q \\ &\geq (1 - \lambda - \gamma)\, c + (1 - \mu - \nu)\, p - \eta g - \xi q \\ &\geq 0; \end{aligned}$$

2) if $-\eta g - \xi q \leq 0$,

$$\begin{aligned} &-\eta g - \xi q \leq 0 \\ \Leftrightarrow\ &(-\eta \cos d_y - \xi \cos d_x) \sin d_z \leq 0 \\ \Leftrightarrow\ &(-\eta \cos d_y - \xi \cos d_x) \sin d_y \leq (-\eta \cos d_y - \xi \cos d_x) \sin d_z \\ \Leftrightarrow\ &-\eta h - \xi p \leq -\eta g - \xi q \end{aligned}$$

where $h = \cos d_y \sin d_y$.

We also have the relation:

$$a \geq \left\{ \begin{array}{c} b \\ c \end{array} \right\} \geq h \geq p \geq 0.$$

So we get

$$\begin{aligned} & a + b - f_2 \\ = & \left(1-\lambda\right) a + \left(1-\mu\right) b - \gamma c - \nu p - \eta g - \xi q \\ \geq & \left(1-\lambda-\gamma\right) c + \left(1-\mu\right) b - \nu p - \eta h - \xi p \\ \geq & \left(2-\lambda-\gamma-\mu-\eta\right) h - \nu p - \xi p \\ \geq & \left(2-\lambda-\gamma-\mu-\eta-\nu-\xi\right) p \\ \geq & \, 0. \end{aligned}$$

Therefore we conclude from above that $a+b \geq f_2$, where the equality is satisfied iff $\theta = 0, \pi$, $\alpha = \pi$, $\omega = \frac{\pi}{2}$ (that is $T^z = -1$).

Let $f' = f_{1max} + f_2 \geq f$. Because $f_{1max}(\omega_1) < f_{1max}(\omega_2)$ when $0 \leq \omega_2 < \omega_1 \leq \pi$, and $f_2 \leq f\left(\theta = 0, \pi\, \omega = \frac{\pi}{2}\right)$, we can get the conclusion that f arrives at its maximum in the domain $0 \leq \omega \leq \pi/2$.

Proposition 3. $f_2 \leq (a+b)\sin\omega$, *when* $0 \leq \omega \leq \frac{\pi}{2}$.

From Eq.(C13) and Eq.(C14), we have the inequality:

$$\lambda + \mu + \gamma + \eta + \nu + \xi \leq 2\sin\omega, \text{ if } 0 \leq \omega \leq \frac{\pi}{2}$$

and

$$\lambda + \mu + \gamma + \eta \leq 2\sin\omega, \text{ if } 0 \leq \omega \leq \frac{\pi}{2}.$$

Also we have the relations:

$$\begin{aligned} \eta + \xi &= T^y S_z^x - T^x S_z^y \\ &= 2\sin\frac{\omega}{2}\sin\alpha\sin\theta\left[\cos\frac{\omega}{2}\cos\left(\phi-\beta\right) - \sin\frac{\omega}{2}\cos\theta\sin\left(\phi-\beta\right)\right] \\ &\leq 2\sin\frac{\omega}{2}\sin\theta\sqrt{\cos^2\frac{\omega}{2} + \sin^2\frac{\omega}{2}\cos^2\theta} \\ &= 2\sin\frac{\omega}{2}\sqrt{-\left(\sin\frac{\omega}{2}\sin^2\theta - \frac{1}{2\sin\frac{\omega}{2}}\right)^2 + \left(\frac{1}{2\sin\frac{\omega}{2}}\right)^2} \end{aligned}$$

Then if $\sin^2\theta = \frac{1}{2\sin^2\frac{\omega}{2}} \leq 1$, then $\eta + \xi \leq 1$. If $\sin^2\frac{\omega}{2} \leq \frac{1}{2}$, i.e., $\omega \leq \frac{\pi}{2}$, then $\eta + \xi \leq \sin\omega$.

Similarly, when $0 \leq \omega \leq \frac{\pi}{2}$, we have

$$\begin{aligned} \lambda + \gamma &\leq \sin\omega, \\ \nu + \mu &\leq \sin\omega, \end{aligned}$$

Therefore we can prove this proposition using the method used in the proof of proposition (2) just replacing 1 and 2 with $\sin\omega$ and $2\sin\omega$ respectively.

3. Conclusion

Therefore

$$\begin{aligned} f &\leq f' \leq f_{1max} + (a+b)\sin\omega \\ &= \sin d_x \sin d_y + \left(\sin d_x + \sin d_y\right)\sin\left(\omega + d_z\right) \\ &\leq \sin d_x \sin d_y + \sin d_x + \sin d_y, \end{aligned}$$

the equality holds when $\alpha = k\pi$, $\theta = (k+1)\pi$, and $\omega = \frac{\pi}{2} - d_z$.

[1] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Phys. Rev. Lett. **70**, 1895 (1993).

[2] J. I. Cirac, P. Zoller, H. J. Kimble, and H. Mabuchi, Phys. Rev. Lett. **78**, 3221 (1997).

[3] S. Bose, Phys. Rev. Lett. **91**, 207901 (2003).

[4] M. Christandl, N. Datta, A. Ekert, and A. J. Landahl, Phys. Rev. Lett. **92**, 187902 (2004).

[5] T. Shi, Y. Li, Z. Song, and C.-P. Sun, Phys. Rev. A **71**, 032309 (2005).

[6] A. Kay, Phys. Rev. Lett. **98**, 010501 (2007).

[7] C. Di Franco, M. Paternostro, and M. S. Kim, Phys. Rev. Lett. **101**, 230502 (2008).

[8] N. Y. Yao, L. Jiang, A. V. Gorshkov, Z.-X. Gong, A. Zhai, L.-M. Duan, and M. D. Lukin, Phys. Rev. Lett. **106**, 040505 (2011).

[9] C. Godsil, S. Kirkland, S. Severini, and J. Smith, Phys. Rev. Lett. **109**, 050502 (2012).

[10] D. Collins, N. Linden, and S. Popescu, Phys. Rev. A **64**, 032302 (2001).

[11] K. Hammerer, G. Vidal, and J. I. Cirac, Phys. Rev. A **66**, 062321 (2002).

[12] P. Zanardi, C. Zalka, and L. Faoro, Phys. Rev. A **62**, 030301 (2000).

[13] B. Kraus and J. I. Cirac, Phys. Rev. A **63**, 062309 (2001).

[14] M. A. Nielsen, C. M. Dawson, J. L. Dodd, A. Gilchrist, D. Mortimer, T. J. Osborne, M. J. Bremner, A. W. Harrow, and A. Hines, Phys. Rev. A **67**, 052301 (2003).

[15] N. Khaneja, R. Brockett, and S. J. Glaser, Phys. Rev. A **63**, 032308 (2001).

[16] W. Rudin, *Functional Analysis* (McGraw-Hill, 1991).